

## NANDA –Approved Nursing Diagnoses

### Pattern 1: Exchanging

Imbalanced nutrition: more than body requirements	Risk for imbalanced fluid volume
Imbalanced nutrition: less than body requirements	Excess fluid volume
Risk for imbalanced nutrition: More than body requirements	Deficient fluid volume
Readiness for enhanced nutrition	Risk for deficient fluid volume
Risk for infection	Readiness for enhanced fluid balance
Risk for imbalanced body temperature	Decreased cardiac output
Hypothermia	Impaired gas exchange
Hyperthermia	Ineffective airway clearance
Ineffective thermoregulation	Ineffective breathing pattern
Autonomic dysreflexia	Impaired spontaneous ventilation
Risk for autonomic dysreflexia	Dysfunctional ventilatory weaning response
Constipation	Risk for injury
Perceived constipation	Risk for falls
Diarrhea	Risk for suffocation
Bowel incontinence	Risk for poisoning
Risk for constipation	Risk for trauma
Impaired urinary elimination	Risk for aspiration
Stress urinary incontinence	Risk for sudden infant death syndrome
Reflex urinary incontinence	Risk for disuse syndrome
Urge urinary incontinence	Latex allergy response
Functional urinary incontinence	Risk for latex allergy response
Total urinary incontinence	Ineffective protection
Risk for urge urinary incontinence	Impaired tissue integrity
Urinary retention	Impaired oral mucous membrane
Readiness for enhanced urinary elimination	Impaired skin integrity
Ineffective tissue perfusion(specify type: renal, cerebral, cardiopulmonary, gastrointestinal, peripheral)	Risk for impaired skin integrity
	Impaired dentition
	Decreased intracranial adaptive capacity
	Disturbed energy field

### Pattern 2: Communicating

Impaired verbal communication`	Readiness for enhanced communication
--------------------------------	--------------------------------------

### *Pattern 3: Relating*

Impaired social interaction	Sexual dysfunction
Social isolation	Interrupted family processes
Risk for loneliness	Readiness for enhanced family processes
Ineffective role performance	Caregiver role strain
Impaired parenting	Risk for caregiver role strain
Risk for impaired parenting	Dysfunctional family process: Alcoholism
Readiness for enhanced parenting	Parental role conflict
Risk for impaired parent/infant/child attachment	Ineffective sexuality patterns

#### ***Pattern 4: Valuing***

Spiritual distress  
Risk for spiritual distress

Readiness for enhanced spiritual well-being

#### ***Pattern 5: Choosing***

Ineffective coping  
Readiness for enhanced coping  
Impaired adjustment  
Defensive coping  
Ineffective denial  
Disabled family coping  
Compromised family coping  
Readiness for enhanced family coping  
Readiness for enhanced community coping  
Ineffective community coping  
Ineffective therapeutic regimen management

Readiness for enhanced management of therapeutic regimen  
Noncompliance (specify)  
Ineffective family therapeutic regimen management  
Ineffective community therapeutic regimen management  
Effective therapeutic regimen management  
Decisional conflict (specify)  
  
Health-seeking behaviors (specify)

#### ***Pattern 6: Moving***

Impaired physical mobility  
Risk for peripheral neurovascular dysfunction  
Risk for perioperative positioning injury  
Impaired walking  
Impaired wheelchair mobility  
Impaired transfer ability  
Impaired bed mobility  
Activity intolerance  
Fatigue  
Risk for activity intolerance  
Disturbed sleep pattern  
Sleep deprivation  
Readiness for enhanced sleep  
Deficient diversional activity  
Impaired home maintenance management  
Ineffective health maintenance  
Delayed surgical recovery  
Adult failure to thrive

Feeding self-care deficit  
Impaired swallowing  
Ineffective breastfeeding  
Interrupted breastfeeding  
Effective breastfeeding  
Ineffective infant feeding pattern  
Bathing/hygiene self-care deficit  
Dressing/grooming self-care deficit  
Toileting self-care deficit  
Delayed growth and development  
Risk for delayed development  
Risk for disproportionate growth  
Relocation stress syndrome  
Risk for relocation stress syndrome  
Risk for disorganized infant behavior  
Disorganized infant behavior  
Readiness for enhanced organized infant behavior

***Pattern 7: Perceiving***

Disturbed body image  
Chronic low self-esteem  
Situational low self-esteem  
Risk for situational low self-esteem  
Readiness for enhanced self-concept  
Powerlessness  
Risk for powerlessness

Disturbed personal identity  
Disturbed sensory perception (specify:  
visual, auditory, kinesthetic, gustatory,  
tactile, olfactory)  
Unilateral neglect  
Hopelessness

***Pattern 8: Knowing***

Deficient knowledge (specify)  
Readiness for enhanced knowledge  
(specify)  
Impaired environmental interpretation  
syndrome

Acute confusion  
Chronic confusion  
Disturbed thought processes  
Impaired memory  
Wandering

***Pattern 9: Feeling***

Acute pain  
Chronic pain  
Nausea  
Dysfunctional grieving  
Anticipatory grieving  
Chronic sorrow  
Risk for other-directed violence  
Self-mutilation  
Risk for self-mutilation  
Risk for self-directed violence

Risk for suicide  
Post-trauma syndrome  
Rape-trauma syndrome  
Rape-trauma syndrome: compound  
reaction  
Rape-trauma syndrome: silent reaction  
Risk for post-trauma syndrome  
Anxiety  
Death anxiety  
Fear